

Two-loop Higgs mass in supersymmetric Kaluza-Klein theories *

A. Delgado, G. v. Gersdorff and M. Quirós

*Instituto de Estructura de la Materia (CSIC), Serrano 123,
E-28006-Madrid, Spain.*

Abstract

We perform an explicit two-loop computation of the Higgs mass in a five-dimensional supersymmetric theory compactified on the orbifold $S^1/\mathbb{Z}_2$, with superpotential localized on a fixed-point brane and supersymmetry broken in the bulk of the extra dimension by Scherk-Schwarz boundary conditions. At one-loop, the Higgs mass is finite and proportional to $1/R^2$, R being the radius of the extra dimension. The two-loop corrections contain finite ($\propto 1/R^2$) and linearly divergent ($\propto \Lambda/R$) terms, where Λ is the cutoff of the five-dimensional theory. The divergent terms are cancelled by diagrams containing one-loop wave-function renormalization counterterms, i.e. by linear threshold effects of Yukawa (and gauge) couplings, at the scale $1/R$. As a result there is no counterterm for the Higgs mass, which is finite and calculable to the considered order.

July 2001

*Work supported in part by CICYT, Spain, under contract AEN98-0816, and by EU under contracts HPRN-CT-2000-00152 and HPRN-CT-2000-00148.

1 Introduction

There has been recently a lot of effort in order to provide an explanation for electroweak symmetry breaking (EWSB) by means of supersymmetric theories with one compact (radius R) extra dimension [1]- [8]. Such theories arise naturally as low energy limits of string theories [9] and are to be understood as effective theories, when all heavy string modes are integrated out, with an ultraviolet (UV) cutoff Λ of the order of the string scale M_s .

A common claim of these models is that electroweak symmetry breaking (EWSB) is triggered by radiative corrections which are insensitive to the scale Λ . Some one loop calculations of the effective potential, or the Higgs mass, have been performed in order to establish this result in field [1]- [8], and string [10] theory, and also some general arguments have been given [5, 11–13].

One uses typically the so called Kaluza-Klein (KK) regularization which exploits the fact that sums over KK modes (integration over the fifth momentum) turn out to be finite (this is already known from finite temperature field theories). Since the summation goes over modes with masses far beyond the cutoff of the effective theory, there have been some doubts on the validity of such a regularization [14]. However it has been shown that several cutoffs which respect all symmetries of the theory lead to a Λ -insensitive result at one-loop, which equals the one obtained in the KK regularization [15, 16]. This is a very interesting conclusion since it means that EWSB does not depend on the microscopic theory at all and we have an effective theory with an unexpected amount of predictability. However, since no rigorous proof has yet been given which guarantees the independence of EWSB on the high energy theory, it seems worthwhile to explicitly check this in a two loop calculation.

In this article we will perform a two-loop calculation of the Higgs mass from the top/stop sector of the model presented in Ref. [8]. It displays the way how supersymmetric cancellations survive the soft breaking. There remains a linear sensitivity to the cutoff which is however entirely absorbed by the supersymmetric running of the Yukawa coupling ¹. As a consequence there is no mass counterterm to the considered order and the Higgs mass is calculable and finite. These features would remain in the two-loop calculation of the Higgs mass when the gauge sector is included.

The outline of this paper is as follows. In section 2 we will summarize the main features of the five-dimensional (5D) supersymmetric model that will be used in the two-loop calculation. In particular the four dimensional lagrangian, after integration over the fifth dimension, with all relevant interactions will be explicitly provided. The corresponding Feynman rules can be easily deduced from the Lagrangian. In section 3 we review, for completeness, the diagrammatic calculation of the one-loop Higgs mass, and the two-loop contribution is provided in section 4, where the relevant one-loop counterterms due to the wave-function renormalization of the superfields, and their contribution to the two-loop Higgs mass, is computed. Our conclusions are drawn in section 5.

¹In the supersymmetric limit there is only a wave-function renormalization for the fields which is absorbed by the running of the Yukawa coupling. This running is of course linear for $D = 5$.

2 The model

The model we will use for the calculation is the one of Ref. [8]. The 5D space-time is compactified on the orbifold $\mathcal{M} \times S^1/\mathbb{Z}_2$ which has two fixed points at $y = 0, \ell$ ($\ell = \pi R$ is the segment length) where two 3-branes are located. There are two types of fields: those living in the bulk of the extra dimension (U -states), organized in $N = 2$ supermultiplets, and those living on the branes and localized at the fixed points (T -states), organized in $N = 1$ supermultiplets. U -states are: gauge fields, which appear in $N = 2$ vector multiplets $\mathbb{V} = (V_\mu, \lambda_1; \Sigma + iV_5, \lambda_2)$ and transform under the adjoint representation of the $SU(3) \times SU(2)_L \times U(1)_Y$, and $SU(2)_L$ singlets, which appear in $N = 2$ hypermultiplets, $\mathbb{U} = (U, u; U', u')$, $\mathbb{D} = (D, d; D', d')$ and $\mathbb{E} = (E, e; E', e')$. Even and odd $N = 1$ superfields under $\mathbb{Z}_2$ are separated by a semicolon as (*even*; *odd*). T -states localized at the $y = 0$ brane are the $SU(2)_L$ doublets, $H_{1,2} = (H_{1,2}, h_{1,2})$, $Q = (Q, q)$ and $L = (L, \ell)$ ². This field assignment allows for a superpotential involving the top/stop sector as

$$W = h_t^{(5)} Q H_2 U \delta(y) \quad (2.1)$$

which triggers EWSB at one-loop as was shown in Ref. [8]. In the rest of this paper we will consider the leading contribution from the top-stop sector and neglect, in explicit calculations, the contributions from gauge couplings. The latter can be easily included and will not modify qualitatively the result of this paper.

Let us express the top/stop sector of the theory in terms of four-dimensional $N = 1$ superfields, using the formalism of Ref. [17].

$$\begin{aligned} \mathcal{L} = & \left(U^\dagger U + U'^\dagger U' + Q^\dagger Q \delta(y) + H_2^\dagger H_2 \delta(y) \right)_{\theta\theta\bar{\theta}\bar{\theta}} \\ & + \left(U \partial_y U' + h_t^{(5)} U Q H_2 \delta(y) + \text{h.c.} \right)_{\theta\theta} \end{aligned} \quad (2.2)$$

The superpotential term is given by the second expression and depends on the chiral superfields Q , H_2 , U and $\partial_y U'$. The inclusion of the derivative term, on top of the superpotential given in (2.1), is pure convention and it is motivated by the fact that, from a four-dimensional viewpoint, it constitutes a mass term for the modes. Notice that we have not included a $\mu H_2 H_1 \delta(y)$ term in the superpotential. The μ parameter is a supersymmetric mass and thus its presence is completely decoupled from the actual calculation of the possible soft supersymmetry Higgs mass counterterms, while it makes the calculation more involved. For that reason we have decided to work in the (phenomenologically unrealistic) case $\mu = 0$. In section 5 we will make some comments about the impact, in the present calculation, of the μ -term. We can anticipate, not unexpectedly, that the final result (in fact the absence of a mass counterterm) will be unchanged by the presence of the μ -term.

²We are using the same notation, upper case symbols, for the $N = 1$ chiral superfields and their scalar components, while lower case symbols are reserved for the fermionic components, e.g. $U = (U, u)$.

In terms of component fields we have

$$\begin{aligned}
\mathcal{L} = & \partial_\mu U^\dagger \partial^\mu U + \partial_\mu U'^\dagger \partial^\mu U' + \partial_\mu Q^\dagger \partial^\mu Q \delta(y) + \partial_\mu H_2^\dagger \partial^\mu H_2 \delta(y) \\
& + i\bar{u}\bar{\sigma}^\mu \partial_\mu u + i\bar{u}'\bar{\sigma}^\mu \partial_\mu u' + i\bar{q}\bar{\sigma}^\mu \partial_\mu q \delta(y) + i\bar{h}_2\bar{\sigma}^\mu \partial_\mu h_2 \delta(y) \\
& - |F_U|^2 - |F_{U'}|^2 - |F_Q|^2 \delta(y) - |F_{H_2}|^2 \delta(y) \\
& + \frac{\partial W}{\partial U} F_U + \frac{\partial W}{\partial U'} F_{U'} + \frac{\partial W}{\partial Q} F_Q + \frac{\partial W}{\partial H_2} F_{H_2} \\
& - \frac{\partial^2 W}{\partial U \partial U'} uu' - \frac{\partial^2 W}{\partial Q \partial U} qu - \frac{\partial^2 W}{\partial Q \partial H_2} qh_2 - \frac{\partial^2 W}{\partial H_2 \partial U} h_2 u + \text{h.c.}
\end{aligned} \tag{2.3}$$

The auxilliary fields are computed to be:

$$F_U^\dagger = -\frac{\partial W}{\partial U} = -h_t^{(5)} \delta(y) Q H_2 - \partial_y U' \tag{2.4}$$

$$F_{U'}^\dagger = -\frac{\partial W}{\partial U'} = \partial_y U \tag{2.5}$$

$$\delta(y) F_Q^\dagger = -\frac{\partial W}{\partial Q} = -h_t^{(5)} \delta(y) U H_2 \tag{2.6}$$

$$\delta(y) F_{H_2}^\dagger = -\frac{\partial W}{\partial H_2} = -h_t^{(5)} \delta(y) U Q \tag{2.7}$$

The interaction and mass terms become

$$\begin{aligned}
\mathcal{L}_{int} = & - \left| h_t^{(5)} \delta(y) Q H_2 + \partial_y U' \right|^2 - |\partial_y U|^2 - \delta(y) \left| h_t^{(5)} U H_2 \right|^2 - \delta(y) \left| h_t^{(5)} U Q \right|^2 \\
& - u \partial_y u' - h_t^{(5)} \delta(y) H_2 q u - h_t^{(5)} \delta(y) U q h_2 - h_t^{(5)} \delta(y) Q h_2 u + \text{h.c.}
\end{aligned} \tag{2.8}$$

Supersymmetry breaking is performed using the Scherk-Schwarz (SS) [18] boundary conditions based on the subgroup $U(1)_R$ which survives after the orbifold $S^1/\mathbb{Z}_2$ action. Using the R -symmetry $U(1)_R$ with parameter ω to impose different boundary conditions for bosons and fermions inside the 5D $N = 1$ multiplets, one obtains for the n th KK mode of gauge bosons and chiral fermions living in the bulk the compactification mass n/R , while for the n th KK mode of gauginos and supersymmetric partners of chiral fermions living in the bulk, the mass $(n + \omega)/R$. Of course, the chiral multiplets localized on the brane do not acquire any tree-level supersymmetry breaking mass.

After SS-breaking, the bulk hypermultiplets have the following Fourier expansions:

$$u = \frac{1}{\sqrt{\pi R}} \sum_{k=-\infty}^{\infty} \cos(ky/R) u^{(k)} \tag{2.9}$$

$$u' = \frac{1}{\sqrt{\pi R}} \sum_{k=-\infty}^{\infty} \sin(ky/R) u^{(k)} \tag{2.10}$$

$$U = \frac{1}{\sqrt{\pi R}} \sum_{k=-\infty}^{\infty} \cos((k + \omega)y/R) U^{(k)} \tag{2.11}$$

$$U' = \frac{1}{\sqrt{\pi R}} \sum_{k=-\infty}^{\infty} \sin((k + \omega)y/R) U^{(k)} \tag{2.12}$$

where we chose to combine the two half-towers to give a single tower.

Plugging these expressions into the Lagrangian and performing the y -integration from 0 to πR we get

$$\begin{aligned}
\mathcal{L} = & \sum_k \left\{ \partial_\mu U^{(k)\dagger} \partial^\mu U^{(k)} - (m_k^B)^2 U^{(k)\dagger} U^{(k)} \right\} + \partial_\mu Q^\dagger \partial^\mu Q + \partial_\mu H_2^\dagger \partial^\mu H_2 \\
& + \sum_k \left\{ i \bar{u}^{(k)} \bar{\sigma}^\mu \partial_\mu u^{(k)} - \frac{1}{2} m_k^F u^{(k)} u^{(k)} - \frac{1}{2} m_k^F \bar{u}^{(k)} \bar{u}^{(k)} \right\} + i \bar{q} \bar{\sigma}^\mu \partial_\mu q + i \bar{h}_2 \bar{\sigma}^\mu \partial_\mu h_2 \\
& - h_t^2 \sum_k Q^\dagger Q H_2^\dagger H_2 - h_t^2 \sum_{k,l} \left\{ U^{(k)\dagger} U^{(l)} H_2^\dagger H_2 + U^{(k)\dagger} U^{(l)} Q^\dagger Q \right\} \\
& - h_t \sum_k \left\{ H_2 q u^{(k)} + U^{(k)} q h_2 + Q h_2 u^{(k)} + m_k^B U^{(k)\dagger} Q H_2 + \text{h.c.} \right\}
\end{aligned} \tag{2.13}$$

where we have defined $h_t^2 = h_t^{(5)2}/\pi R$, $m_k^F = k/R$ and $m_k^B = (k + \omega)/R$ and made use of the “representation” $\delta(0) = (\pi R)^{-1} \sum_{-\infty}^{\infty} 1$. Had we kept two separate half towers of fermions one could combine the nonzero modes to give a Dirac fermion.

3 One-loop contribution to the Higgs mass

In this section we compute, for the sake of completeness and to fix some notations, the one-loop contribution to the Higgs mass. We define the propagators of the bulk fields with end points at $y = 0$ as:

$$D^\omega(p) = \sum_{n=-\infty}^{\infty} \frac{1}{p^2 - \left(\frac{n+\omega}{R}\right)^2} = \frac{\pi R}{2p} (\cot(\pi(pR + \omega)) + \cot(\pi(pR - \omega))) \tag{3.1}$$

$$\tilde{D}^\omega(p) = \sum_{n=-\infty}^{\infty} \frac{\frac{n+\omega}{R}}{p^2 - \left(\frac{n+\omega}{R}\right)^2} = -\frac{\pi R}{2} (\cot(\pi(pR + \omega)) - \cot(\pi(pR - \omega))) \tag{3.2}$$

For Euclidean momenta ($p \rightarrow ip$)

$$\begin{aligned}
D_E^\omega(p) &= -\frac{\pi R}{2p} (\coth(\pi(pR + i\omega)) + \coth(\pi(pR - i\omega))) \\
\tilde{D}_E^\omega(p) &= \frac{\pi R}{2i} (\coth(\pi(pR + i\omega)) - \coth(\pi(pR - i\omega)))
\end{aligned} \tag{3.3}$$

the propagators have the asymptotic expansions

$$D_E^\omega(p) = -\frac{\pi R}{p} (1 + 2 e^{-2\pi R p} \cos(2\pi\omega)) \tag{3.4}$$

$$\tilde{D}_E^\omega(p) = 2\pi R e^{-2\pi R p} \sin(2\pi\omega) \tag{3.5}$$

From Eq. (3.4) we can see that $D_E^\omega(p)$ has a leading term $\sim 1/p$ which is independent of ω while the remaining part decays exponentially. This reflects the soft nature of the supersymmetry breaking by the SS mechanism.

At one loop there are four diagrams which are shown in Fig. 1. They contribute [2] to the Higgs mass as

$$\begin{aligned}
m_H^2 &= iN_c h_t^2 \int \frac{d^4 q}{(2\pi)^4} \left\{ -2D^0(q) + \sum_k \frac{(m_k^B)^2}{q^2 (q^2 - (m_k^B)^2)} + \sum_k \frac{1}{q^2} + D^\omega(q) \right\} \\
&= -\frac{N_c h_t^2}{4\pi^2} \int_0^\infty dq \{ D_E^\omega(q) - D_E^0(q) \}
\end{aligned} \tag{3.6}$$




Figure 1: The diagrams contributing to the Higgs mass at one loop.

The difference $D_E^\omega(q) - D_E^0(q)$ decays exponentially (as can be seen from Eq. (3.4)) so the result is finite. The integral has been evaluated in Refs. [1, 2, 8]:

$$m_H^2 = \frac{N_c h_t^2}{16\pi^4 R^2} [Li_3(e^{-2i\pi\omega}) + Li_3(e^{2i\pi\omega}) - 2\zeta(3)] \tag{3.7}$$

There is a cancellation of tree level infinities between the second and third diagrams. This cancellation always occurs according to the diagrammatical rule

$$\begin{array}{c} H_2 \\ \diagdown \end{array} \begin{array}{c} \text{---} \end{array} \begin{array}{c} \diagup \\ H_2 \end{array} + \begin{array}{c} H_2 \\ \diagdown \end{array} \begin{array}{c} \text{---} \end{array} \begin{array}{c} \diagup \\ H_2 \end{array} = -i h_t^2 q^2 D^\omega(q) \tag{3.8}$$

The corresponding rule will be to drop all diagrams containing $\delta(0)$ and replace in the remaining ones

$$\sum_k \frac{m_k^2}{q^2 - m_k^2} \rightarrow q^2 D^\omega(q) \tag{3.9}$$

4 Two-loop contribution to the Higgs mass

In this section we compute the leading two-loop contributions to the Higgs mass coming from the top/stop sector, as we did at one-loop in section 3, i.e. we consider $\mathcal{O}(h_t^4)$

corrections. We use for that the Lagrangian (2.13). Inclusion of either gauge couplings or a $\mu \neq 0$ term will not alter the qualitative conclusions of our analysis. We will comment more on that in section 5.

In order to display explicitly the various cancellations we group the genuine two-loop diagrams as shown in figures 2 to 8, and the one-loop diagrams with counterterm insertions in figures 9 and 10. In all cases external legs correspond to H_2 fields.

4.1 Two-loop diagrams

First there are the three diagrams shown in figure 2.




Figure 2: Diagrams which vanish identically.

The third one does actually not exist since there is no corresponding contraction providing the corresponding propagators (fields are complex). In all diagrams containing fermions, the terms proportional to the masses cancel, since the masses are zero for q and h and for the case of the $u^{(k)}$ cancellation occurs due to symmetric summation over an odd function of k . For this reason diagrams **0a** and **0b** are zero when summed over.

Contractions are done using Weyl fermions. The relevant propagators are:

$$\langle T(\psi_\alpha \bar{\psi}_{\dot{\alpha}}) \rangle(p) = \frac{p_m \sigma_{\alpha\dot{\alpha}}^m}{p^2 - m^2} \quad (4.1)$$

$$\langle T(\psi^\alpha \bar{\psi}^{\dot{\alpha}}) \rangle(p) = \frac{p^m \bar{\sigma}_m^{\dot{\alpha}\alpha}}{p^2 - m^2} \quad (4.2)$$

and the contraction rules:

$$\begin{aligned} \text{Tr } \sigma^m \bar{\sigma}^n &= -2\eta^{mn} \\ \text{Tr } \sigma^m \bar{\sigma}^n \sigma^r \bar{\sigma}^s &= -2(\eta^{mn}\eta^{rs} - \eta^{mr}\eta^{ns} + \eta^{ms}\eta^{nr}) \end{aligned} \quad (4.3)$$

are used.

There will be two kinds of cancellation between bosons and fermions:

- If a loop involves only particles which are localized on the brane, there will be the usual supersymmetric cancellations of quadratic divergences. Supersymmetry breaking does not affect these particles (both fermions and bosons remain massless).
- If a loop involves particles moving in the bulk, one expects cubic divergences by power counting. In the supersymmetric theory ($\omega = 0$) these terms are cancelled

exactly by the sfermion. For $\omega \neq 0$ fermions and bosons get different masses. However after summation over the modes (integration over the loop fifth momentum) the ω -dependence lies in the propagators D^ω and D^0 whose difference is exponentially suppressed for large euclidian four-momentum (see Eq. (3.4)). So the cancellation is no longer exact but leaves over a UV-finite piece.

We use in the following the definitions $k = p + q$, $l = p - q$ and the identities

$$\begin{aligned} 2p \cdot q &= k^2 - p^2 - q^2 \\ &= -l^2 + p^2 + q^2 \\ 2p \cdot k &= -q^2 + p^2 + k^2 \\ 2q \cdot k &= -p^2 + q^2 + k^2 \end{aligned} \tag{4.4}$$

to eliminate the scalar products and express all integrands in terms of p^2 , q^2 and k^2 . We will suppress the integrations and the common factor of h_t^4 . The diagrams of the first group are shown in figure 3.




Figure 3: Two-loop diagrams computed in Eqs. (4.5)-(4.7). Diagrams **1c** and **1d** have the same value.

The calculation of the diagrams in Fig. 3 leads to expressions $d_1 = d_{1a} + d_{1b} + d_{1c} + d_{1d}$.

$$\begin{aligned} d_{1a} &= -2iN_c [D^\omega(q)]^2 \frac{p \cdot k}{p^2 k^2} = -iN_c [D^\omega(q)]^2 \left(-\frac{q^2}{p^2 k^2} + \frac{1}{p^2} + \frac{1}{k^2} \right) \\ &= -iN_c [D^\omega(q)]^2 \left(-\frac{q^2}{p^2 k^2} + \frac{2}{p^2} \right) \end{aligned} \tag{4.5}$$

The possibility of the renaming $k \rightarrow p$ becomes obvious if one writes the integration over k explicitly as $\int d^4k \delta(p + q - k)$ and observes that the sharp cutoff has to be introduced symmetrically for all three momenta.

$$d_{1b} = iN_c \left[\tilde{D}^\omega(q) \right]^2 \frac{1}{k^2 p^2} \tag{4.6}$$

$$d_{1c} = d_{1d} = iN_c [D^\omega(q)]^2 \frac{1}{p^2} \tag{4.7}$$

Assembling the diagrams of Fig. 3 we get

$$d_1 = iN_c [D^\omega(q)]^2 \frac{q^2}{p^2 k^2} + iN_c [\tilde{D}^\omega(q)]^2 \frac{1}{k^2 p^2} \quad (4.8)$$

Note the supersymmetric cancellation of the quadratic divergences in the p -integration. The cancellation involves only loops in the brane and it is identical to the one that would occur in an unbroken theory.

Now consider the class of diagrams of figure 4.




Figure 4: Two-loop diagrams computed in Eqs.(4.9)-(4.12). Diagrams **2b** and **2c** are equal. The last three diagrams combine according to the rule (3.8). Note there is indeed a combinatorial factor of 2 for diagram **2e**, as needed.

$$d_{2a} = -2iN_c [\tilde{D}^\omega(q)]^2 \frac{p \cdot k}{q^2 p^2 k^2} = -iN_c [\tilde{D}^\omega(q)]^2 \left(\frac{1}{q^2 p^2} - \frac{1}{p^2 k^2} + \frac{1}{q^2 k^2} \right) \quad (4.9)$$

$$= -iN_c [\tilde{D}^\omega(q)]^2 \left(\frac{2}{q^2 p^2} - \frac{1}{p^2 k^2} \right) \quad (4.10)$$

$$d_{2b} = d_{2c} = iN_c [\tilde{D}^\omega(q)]^2 \frac{1}{q^2 p^2} \quad (4.11)$$

$$d_{2d} + d_{2e} + d_{2f} = iN_c [D^\omega(q)]^2 \frac{q^2}{p^2 k^2} \quad (4.12)$$

Adding these we get a cancellation of the quadratic divergence in the p -integration:

$$d_2 = iN_c [D^\omega(q)]^2 \frac{q^2}{p^2 k^2} + iN_c [\tilde{D}^\omega(q)]^2 \frac{1}{k^2 p^2} \quad (4.13)$$

Again this cancellation is the unmodified supersymmetric cancellation.

The next contribution comes from correction of the u propagator (see figure 5).




Figure 5: Two-loop diagrams computed in Eq. (4.14).

$$\begin{aligned}
 d_{3a} = d_{3b} &= -2iN_c [D^0(q)]^2 \frac{q \cdot p q^2}{p^2 q^2 l^2} = -iN_c [D^0(q)]^2 \left(-\frac{1}{p^2} + \frac{1}{k^2} + \frac{q^2}{p^2 k^2} \right) \\
 &= -iN_c [D^0(q)]^2 \frac{q^2}{p^2 k^2}
 \end{aligned} \tag{4.14}$$

Adding d_1 to d_3 we get

$$d_1 + d_2 + d_3 = 2iN_c \left([D^\omega(q)]^2 - [D^0(q)]^2 \right) \frac{q^2}{p^2 k^2} + 2iN_c [\tilde{D}^\omega(q)]^2 \frac{1}{k^2 p^2} \tag{4.15}$$

Now the q -integration is completely finite. The remaining logarithmic divergence of the p -integration will be associated to the wave-function renormalization (WFR) of the bulk-field.

Next we consider the diagrams from figure 6.




Figure 6: Two-loop diagrams computed in Eqs. (4.16)-(4.18). They have to be combined according to rule (3.8), i.e. **4a+4b**, **4c+4d**, **4e+4f+4g+4h**.

All of them are infinite by themselves (having $\delta(0)$ vertices or divergent sums) but combining them as below these divergences cancel:

$$d_{4a} + d_{4b} = -2iN_c D^\omega(q) D^0(p) \frac{p \cdot k}{q^2 k^2} = -iN_c D^\omega(q) D^0(p) \left(-\frac{1}{k^2} + \frac{p^2}{q^2 k^2} + \frac{1}{q^2} \right) \tag{4.16}$$

$$d_{4c} + d_{4d} = iN_c D^\omega(q) D^\omega(p) \frac{1}{q^2} \quad (4.17)$$

$$d_{4e} + d_{4f} + d_{4g} + d_{4h} = iN_c D^\omega(q) D^\omega(p) \frac{p^2}{q^2 k^2} \quad (4.18)$$

Summing these up we find a supersymmetric cancellation of a cubic divergence in the integration over p :

$$d_4 = iN_c D^\omega(q) (D^\omega(p) - D^0(p)) \left(\frac{p^2}{q^2 k^2} + \frac{1}{q^2} \right) + iN_c D^\omega(q) D^0(p) \frac{1}{k^2} \quad (4.19)$$

We also have the corrections to the fermionic (q) propagator (see figure 7):

$$d_{5a} = -2iN_c D^\omega(p) D^0(q) \frac{q \cdot k}{q^2 k^2} = -iN_c D^\omega(p) D^0(q) \left(-\frac{p^2}{q^2 k^2} + \frac{1}{k^2} + \frac{1}{q^2} \right) \quad (4.20)$$

$$d_{5b} = -2iN_c D^0(q) D^0(p) \frac{p \cdot q}{q^2 l^2} = -iN_c D^0(q) D^0(p) \left(-\frac{1}{q^2} + \frac{1}{k^2} + \frac{p^2}{q^2 k^2} \right) \quad (4.21)$$

They combine to give a supersymmetric cancellation of the cubic divergence in the integration over p :

$$\begin{aligned} d_5 = & -iN_c D^0(q) (D^\omega(p) - D^0(p)) \left(-\frac{p^2}{q^2 k^2} + \frac{1}{q^2} \right) \\ & - iN_c D^0(q) (D^\omega(p) + D^0(p)) \frac{1}{k^2} \end{aligned} \quad (4.22)$$

Summing up the two contributions we get:

$$\begin{aligned} d_4 + d_5 = & iN_c (D^\omega(q) - D^0(q)) (D^\omega(p) - D^0(p)) \frac{1}{q^2} \\ & + iN_c (D^\omega(q) + D^0(q)) (D^\omega(p) - D^0(p)) \frac{p^2}{q^2 k^2} \\ & + iN_c (D^\omega(q) D^0(p) - D^0(q) (D^\omega(p) + D^0(p))) \frac{1}{k^2} \end{aligned} \quad (4.23)$$

There is a linear divergence in the p -integration coming from the last term. It is associated with the WFR of the matter fields living on the brane and will be cancelled by the usual supersymmetric counterterm.




Figure 7: Two-loop diagrams computed in Eqs. (4.20)-(4.21).

To make the q -integration finite we look at the three remaining diagrams (see figure 8)




Figure 8: Last three two-loop contributions, computed in Eqs. (4.24)-(4.26). Diagrams **7** and **8** are UV-finite

$$d_6 = iN_c D^\omega(q) D^\omega(p) \frac{1}{k^2} \quad (4.24)$$

$$d_7 = iN_c^2 \tilde{D}^\omega(p) \tilde{D}^\omega(q) \frac{1}{q^2 p^2} \quad (4.25)$$

$$d_8 = 2iN_c \tilde{D}^\omega(p) \tilde{D}^\omega(q) \frac{1}{k^2 p^2} \quad (4.26)$$

d_7 and d_8 are UV finite according to the expansion (3.5), while d_6 is exactly the missing piece in equation (4.23).

We end up with:

$$\begin{aligned} -N_c h_t^4 \int \frac{d^4 p}{(2\pi)^4} \frac{d^4 q}{(2\pi)^4} \left\{ 2 \left([D_E^\omega(q)]^2 - [D_E^0(q)]^2 \right) \frac{q^2}{p^2 k^2} - 2 \left[\tilde{D}_E^\omega(q) \right]^2 \frac{1}{k^2 p^2} \right. \\ + (D_E^\omega(q) - D_E^0(q)) (D_E^\omega(p) - D_E^0(p)) \frac{1}{q^2} \\ + (D_E^\omega(q) + D_E^0(q)) (D_E^\omega(p) - D_E^0(p)) \frac{p^2}{q^2 k^2} \\ + (D_E^\omega(q) - D_E^0(q)) (D_E^\omega(p) + D_E^0(p)) \frac{1}{k^2} \\ \left. - N_c \tilde{D}_E^\omega(p) \tilde{D}_E^\omega(q) \frac{1}{q^2 p^2} - 2 \tilde{D}_E^\omega(p) \tilde{D}_E^\omega(q) \frac{1}{k^2 p^2} \right\} \end{aligned} \quad (4.27)$$

One sees that the only remaining divergences are in the p -integration, more specifically there is the logarithmic divergence in the first line and the linear divergence in the fourth. The expression vanishes for $\omega = 0$.

4.2 The counterterms

As in any (exact or softly broken) supersymmetric theory the superpotential couplings run due to the wave-function renormalization of the superfields. We would like to rescale the fields which couple to the brane in a manifestly supersymmetric way. To this end we

rescale the fields as follows:

$$Q \rightarrow \left(1 - \frac{\delta_Q}{2}\right) Q \quad q \rightarrow \left(1 - \frac{\delta_Q}{2}\right) q \quad (4.28)$$

$$H_2 \rightarrow \left(1 - \frac{\delta_{H_2}}{2}\right) H_2 \quad h_2 \rightarrow \left(1 - \frac{\delta_{H_2}}{2}\right) h_2 \quad (4.29)$$

$$U \rightarrow \left(1 - \frac{\delta_U}{2} \pi R \delta(y)\right) U \quad u \rightarrow \left(1 - \frac{\delta_U}{2} \pi R \delta(y)\right) u \quad (4.30)$$

$$\partial_y U' \rightarrow \left(1 + \frac{\delta_U}{2} \pi R \delta(y)\right) \partial_y U' \quad \partial_y u' \rightarrow \left(1 + \frac{\delta_U}{2} \pi R \delta(y)\right) \partial_y u' \quad (4.31)$$

$\partial_y U$ and U' do not couple to the brane and thus will not receive a brane rescaling. We define also a rescaled (renormalized) coupling as

$$h_t = \left(1 - \frac{\delta_Q}{2} - \frac{\delta_{H_2}}{2} - \frac{\delta_U}{2} \pi R \delta(0)\right) h_t^{bare} \quad (4.32)$$

By inspection of equation (2.2) it is obvious that this leaves the Lagrangian unchanged if expressed in rescaled quantities. The different sign for U and $\partial_y U'$ guarantees that the first term of the superpotential remains unaltered.

The following counterterms now appear in the 4D Lagrangian:

$$\begin{aligned} \mathcal{L}_{CT} = & -\delta_Q \partial_\mu Q^\dagger \partial_\mu Q - \delta_{H_2} \partial_\mu H_2^\dagger \partial_\mu H_2 - \delta_U \pi R (\partial_\mu U^\dagger \partial_\mu U + \partial_y U'^\dagger \partial_y U')_{y=0} \\ & -\delta_Q q \sigma^\mu \partial_\mu \bar{q} - \delta_{H_2} h_2 \sigma^\mu \partial_\mu \bar{h}_2 - \delta_U \pi R (u \sigma^\mu \partial_\mu \bar{u})_{y=0} \\ & -\delta_Q h_t^2 \pi R (U^\dagger U)_{y=0} H_2^\dagger H_2 - \delta_{H_2} h_t^2 \pi R (U^\dagger U)_{y=0} Q^\dagger Q \\ & -\delta_U h_t^2 (\pi R \delta(0))^2 Q^\dagger Q H_2^\dagger H_2 + \delta_U h_t \pi R \delta(0) (Q H_2 (\partial_y U')_{y=0} + \text{h.c.}) \\ = & -\delta_Q \partial_\mu Q^\dagger \partial_\mu Q - \delta_{H_2} \partial_\mu H_2^\dagger \partial_\mu H_2 - \delta_U \sum_{k, l} \partial_\mu U^{(k)\dagger} \partial_\mu U^{(l)} - \delta_U \sum_{k, l} m_k U^{(k)\dagger} m_l U^{(l)} \\ & -\delta_Q q \sigma^\mu \partial_\mu \bar{q} - \delta_{H_2} h_2 \sigma^\mu \partial_\mu \bar{h}_2 - \delta_U \sum_{k, l} u^{(k)} \sigma^\mu \partial_\mu \bar{u}^{(l)} \\ & -\delta_Q h_t^2 \sum_{k, l} U^{(k)\dagger} U^{(l)} H_2^\dagger H_2 - \delta_{H_2} h_t^2 \sum_{k, l} U^{(k)\dagger} U^{(l)} Q^\dagger Q \\ & -\delta_U h_t^2 (\pi R \delta(0))^2 Q^\dagger Q H_2^\dagger H_2 + \delta_U h_t \pi R \delta(0) \left(\sum_k m_k U^{(k)} Q H_2 + \text{h.c.} \right) \end{aligned} \quad (4.33)$$

The values of δ_U and δ_Q are determined by the infinite parts of the one-loop two point functions at nonvanishing momentum q . One finds:

$$\begin{aligned} \langle U_k^\dagger U_l \rangle &= h_t^2 \int \frac{d^4 p}{(2\pi)^4} \frac{m_k^B m_l^B + q^2}{p^2 k^2} \\ &= i h_t^2 (m_k^B m_l^B + q^2) \frac{1}{8\pi^2} \log \left(\frac{\Lambda^2}{\mu^2} \right) + \text{finite} \end{aligned} \quad (4.34)$$

$$\begin{aligned}
\langle Q^\dagger Q \rangle &= h_t^2 \int \frac{d^4 p}{(2\pi)^4} [D^\omega(p) - D^0(p)] \left(1 + \frac{p^2}{k^2}\right) + \frac{q^2}{k^2} D^0(p) \\
&= i h_t^2 q^2 \frac{R\Lambda}{8\pi} + \text{finite}
\end{aligned} \tag{4.35}$$

where μ is an infrared cutoff. The infinities can be absorbed if one chooses

$$\delta_U = h_t^2 \frac{1}{8\pi^2} \log \left(\frac{\Lambda^2}{\mu^2} \right) \tag{4.36}$$

$$\delta_Q = h_t^2 \frac{R\Lambda}{8\pi} \tag{4.37}$$

There are now the additional diagrams shown in figures 9 and 10 (δ_U counterterms are denoted by $\oplus$ and δ_Q counterterms by $\oplus$).

The diagrams in figure 9 contribute:

$$2\delta_Q (D^\omega(q) - D^0(q)) \tag{4.38}$$




Figure 9: Counterterm diagrams, computed in Eq. (4.38), proportional to δ_Q .

The diagrams of figure 10 give (observe the factor of two for the third diagram):

$$\begin{aligned}
&\delta_U \sum_{k,l} \left(\frac{1}{q^2} \frac{m_k}{q^2 - m_k^2} \frac{m_l}{q^2 - m_k^2} (q^2 + m_k m_l) + 2 \frac{1}{q^2} \frac{m_k^2}{q^2 - m_k^2} \right. \\
&\quad \left. + \frac{1}{q^2 - m_k^2} \frac{1}{q^2 - m_k^2} (q^2 + m_k m_l) + \frac{1}{q^2} \right) - 2\delta_U q^2 [D^0(q)]^2 \\
&= \delta_U \sum_{k,l} \frac{1}{q^2} \frac{1}{q^2 - m_k^2} \frac{1}{q^2 - m_k^2} \left((q^2 + m_k m_l) m_k m_l + 2(q^2 - m_l^2) m_k^2 \right. \\
&\quad \left. + q^2(q^2 + m_k m_l) + (q^2 - m_k^2)(q^2 - m_l^2) \right) - 2\delta_U q^2 [D^0(q)]^2 \\
&= 2\delta_U \left(q^2 [D^\omega(q)]^2 - q^2 [D^0(q)]^2 + [\tilde{D}^\omega(q)]^2 \right) \tag{4.39}
\end{aligned}$$




Figure 10: Counterterm diagrams, computed in Eq. (4.39), proportional to δ_U .

So the total counterterm contribution is:

$$-N_c h_t^2 \int \frac{d^4 q}{(2\pi)^4} 2\delta_Q (D_E^\omega(q) - D_E^0(q)) - 2\delta_U \left(q^2 [D_E^\omega(q)]^2 - q^2 [D_E^0(q)]^2 - [\tilde{D}_E^\omega(q)]^2 \right) \quad (4.40)$$

Adding them to the divergent terms of (4.27) we get

$$\begin{aligned} -N_c h_t^2 \int \frac{d^4 q}{(2\pi)^4} \left\{ 2 \left(q^2 [D_E^\omega(q)]^2 - q^2 [D_E^0(q)]^2 - [\tilde{D}_E^\omega(q)]^2 \right) \left(h_t^2 \int \frac{d^4 p}{(2\pi)^4} \frac{1}{k^2 p^2} - \delta_U \right) \right. \\ \left. + (D_E^\omega(q) - D_E^0(q)) \left(h_t^2 \int \frac{d^4 p}{(2\pi)^4} (D_E^\omega(p) + D_E^0(p)) \frac{1}{k^2} + 2\delta_Q \right) \right\} \quad (4.41) \end{aligned}$$

which is finite if the counterterms are chosen according to (4.36, 4.37).

5 Conclusions

In this paper we have made an explicit two-loop calculation of the Higgs mass in the five dimensional model of Ref. [8], where the fifth dimension is compactified on the orbifold $S^1/\mathbb{Z}_2$ and supersymmetry is broken in the bulk by means of Scherk-Schwarz boundary conditions. We have considered only (top) Yukawa couplings in the calculation (h_t) and disregarded gauge couplings (g). We have also considered $\mathcal{O}(\mu^0)$ results, which amounts to disregard the μ -parameter in the superpotential. Our result is that no Higgs mass counterterm is generated at two-loop and the only divergent terms in the two-loop correction can be absorbed by the renormalization group running of h_t between the cutoff Λ and the scale where the theory becomes four dimensional, $1/R$. The latter can be interpreted as threshold effects at $1/R$ when matching the 4D theory below it with the complete KK theory above $1/R$. As a result our explicit calculation shows that the Higgs mass is finite and calculable to the considered order in perturbation theory.

Inclusion of gauge couplings has been done at one-loop, $\mathcal{O}(g^2)$ terms, for an arbitrary gauge group [1, 2]. At two-loop we would have $\mathcal{O}(g^2 h_t^2)$ and $\mathcal{O}(g^4)$ terms. We expect divergent terms which are canceled by supersymmetry or by counterterms from

supersymmetric wave-function renormalization, which can be absorbed by the running of gauge and Yukawa couplings, yielding again a finite result.

Inclusion of the μ -term (a supersymmetric mass) in the superpotential would introduce divergences, even at the one-loop level, that can be interpreted as the supersymmetric running of the tree level Higgs mass terms, while the analysis becomes much more involved since the Higgs mass $\mathcal{M}^2$ becomes a two-by-two matrix. All infinities, in this case, are expected to be absorbed by supersymmetry as well as by the running of gauge and Yukawa couplings, and of the μ -parameter. The resulting absence of Higgs mass counterterms should hold in this case. At one-loop, we have checked that the field rescaling $\delta_{H_2} = N_c \delta_Q$ is indeed sufficient to render $\mathcal{M}^2$ finite.

Acknowledgements

The work of AD was supported by the Spanish Education Office (MEC) under an *FPI* scholarship. The work of GG was supported by the DAAD.

References

- [1] I. Antoniadis, S. Dimopoulos, A. Pomarol and M. Quirós, *Nucl. Phys.* **B554** (1999) 503.
- [2] A. Delgado, A. Pomarol and M. Quirós, *Phys. Rev.* **D60** (1999) 095008.
- [3] R. Barbieri, L.J. Hall and Y. Nomura, *Phys. Rev.* **D63** (2001) 105007.
- [4] R. Barbieri, L.J. Hall and Y. Nomura, [hep-ph/0106190](#).
- [5] R. Barbieri, L.J. Hall and Y. Nomura, [hep-th/0107004](#).
- [6] N. Arkani-Hamed, L.J. Hall, Y. Nomura, D. Smith and N. Weiner, *Nucl. Phys.* **B605** (2001) 81.
- [7] A. Delgado and M. Quirós, *Phys. Lett.* **B484** (2000) 355.
- [8] A. Delgado and M. Quirós, [hep-ph/0103058](#).
- [9] For recent reviews see: I. Antoniadis and K. Benakli, *Int. J. Mod. Phys. A* **15** (2000) 4237; I. Antoniadis, [hep-th/0102202](#).
- [10] I. Antoniadis, K. Benakli and M. Quirós, *Nucl. Phys.* **B583** (2000) 35.
- [11] Y. Nomura, [hep-ph/0105113](#).
- [12] N. Weiner, [hep-ph/0106021](#).
- [13] A. Masiero, C.A. Scrucca, M. Serone and L. Silverstrini, [hep-ph/0107201](#).
- [14] D. Ghilencea and H. P. Nilles, *Phys. Lett.* **B507** (2001) 327.

- [15] A. Delgado, G. von Gersdorff, P. John and M. Quirós, [hep-ph/0104112](#).
- [16] R. Contino and L. Pilo, [hep-ph/0104130](#).
- [17] N. Arkani-Hamed, T. Gregoire and J. Wacker, [hep-th/0101233](#); N. Arkani-Hamed, L. Hall, D. Smith and N. Weiner, *Phys. Rev.* **D63** (2001) 056003.
- [18] J. Scherk and J.H. Schwarz, *Phys. Lett.* **B82** (1979) 60; J. Scherk and J.H. Schwarz, *Nucl. Phys.* **B153** (1979) 61; P. Fayet, *Phys. Lett.* **B159** (1985) 121; P. Fayet, *Nucl. Phys.* **B263** (1986) 649.